# **OpenHyPE: OpenHygrisC Data Processing for Education**

- Project funded by the Federal State of Northrhine-Westphalia (NRW)
- Funded by Ministerium für Landwirtschaft und Verbraucherschutz NRW (MLV)
- Project duration: 15.12.2021 30.06.2023

**U**ZUSAMMENFASSUNG, EINLEITUNG UND ZIELE AUF DEUTSCH

# 0. Abstract

The State Agency for Nature, Environment and Consumer Protection (LANUV) of the federal state of North Rhine-Westphalia (NRW) provides extensive quantitative and qualitative groundwater monitoring data. This contributes to the fulfillment of the European Water Framework Directive as well as the EU INSPIRE Directive for an open interoperable spatial data infrastructure.

NRW operates its own water-related data portal **ELWAS-WEB**, which also provides the groundwater database **HygrisC**. ELWAS and HygrisC are not easy to use and provide only limited exploratory data analysis capabilities to the public.

However, the state publishes much of its groundwater data as an **open data archive** called **OpenHygrisC**, which contains several data tables in **csv format**. In particular, the **big data** in the measurement table, which contains all time series with **more than 3.6 million individual measurements** (table rows), and the table with the spatial coordinates of the groundwater monitoring wells require the use of a **spatially enabled object-relational database management system (Spatial ODBRMS)** and extensive **data engineering** before insertion into the database.

The goal of the **OpenHyPE project** is to develop a first set of **Open Educational Resources (OER)** to train the setup, filling and use of a **geospatial-temporal database** with the OpenHygrisC data. Due to the **graduated level of difficulty**, the project addresses **students** from **secondary schools** as well as **universities** in the state of North Rhine-Westphalia and beyond.

All software products used are **Free and Open Source Software (FOSS)**. The database, which we call **OpenHyPE DB**, is based on **PostgreSQL / PostGIS** and establishes the center of the system for environmental data analysis and presentation. The OER demonstates how the geographic information system **QGIS** as well as **Python programs** in the **JupyterLab development environment** interoperate with the OpenHyPE DB to select, analyze and display the data in the form of time dependent maps or time series. We use Python and Jupyter from the **Anaconda distribution**.

The start-up funding for the OpenHyPE project is used to raise awareness of the **NRW's valuable** open environmental data collection among young people as well as to contribute to interdisciplinary STEM promotion in general and education for sustainable development (ESD) in particular by linking environmental science and computer science.

### **1. Introduction**

### **1.1 Problem Description**

The state of North Rhine-Westphalia (NRW) operates comprehensive and professional measurement networks for the collection of environmental data through the LANUV. As part of Open.NRW and driven by the INSPIRE Directive of the European Union as well as other directives such as the EU Water Framework Directive (WFD), the state of NRW makes extensive data products openly accessible and freely usable on various platforms (Free and Open Data).

The state of NRW is a pioneer in Germany in providing open and (cost)free geodata. These data are a real treasure and form the basis for potentially massive knowledge gains in the field of environmental and nature conservation. Nevertheless, it seems that only a comparatively small group of people really uses this potential. Therefore, the OpenHyPE project has set itself the task of integrating this data stock into university teaching and developing corresponding freely accessible teaching material that can be used not only by students but also, to some extent, by pupils to learn the basics of environmental data processing. The start-up funding will be used to implement the first steps of developing such training material.

We follow the paradigm of "problem based learning": the necessary knowledge and skills are identified and taught based on a concrete socially relevant problem. The solution of the problem identified as significant is the motivation for learning.

At the beginning we want to develop the material on the basis of the problem area "groundwater protection". The Ministry for Environment, Agriculture, Nature and Consumer Protection NRW (MULNV) operates its own water-related data portal called ELWAS-WEB via the "Landesbetrieb Information und Technik Nordrhein-Westfalen" (IT.NRW). Data from the statewide groundwater database HygrisC are also held in this portal. ELWAS and HygrisC offer limited exploratory data analysis capabilities to outsiders. From the point of view of usability engineering, which deals with the user-friendliness of technical systems, improvements are desirable with regard to usability as well as data analysis possibilities, because exploratory data analysis and data mining in particular help to identify structures and relationships between data. ELWAS and HygrisC are therefore only suitable to a limited extent for teaching the basics of environmental data analysis, but they can be used in the classroom as supporting material.

On the portal OpenGeodata.NRW extensive data with spatial reference - also called geodata - are made available, which often have a time reference, such as land use changes or measurement data series on water quality. Excerpts of the HygrisC groundwater database of the state of NRW, published under the name OpenHygrisC, are also located there. These groundwater data can ideally serve as a basis for building one's own environmental database, which the learners can use to learn about concepts of data management and data analysis.

### 1.2 Project Goals

The following components are to be realized:

- Development of OpenHyPE geodatabase based on PostgreSQL/PostGIS to manage spatial and temporal data on groundwater quality and quantity.
- Problem-related free online course material (OER), tutorials, video tutorials, instructions,

#### program code,

using Free and Open Source Software (FOSS):

- Introduction to the State Agency for Nature, Environmental and Consumer Protection (LANUV).
- Introduction to groundwater protection
- $\circ\,$  Introduction to the Geographic Information System QGIS
- $\circ\,$  Introduction to the relational database PostgreSQL and the query language SQL
- Introduction to the geodatabase extension PostGIS
- $\circ\,$  Introduction to the processing of geodata with the programming language Python
- $\circ~$  Installation of the OpenHyPE database management system
- Discussion of the data model and upload of the OpenHygrisC data of the LANUV
- Automatic creation of diagrams on time series of water quality
- Automatic generation of groundwater chemistry maps
- $\circ\,$  Creating simple dashboards with interactive online graphs and maps
- Introduction to data mining (descriptive statistics, searching for correlations)

# 2. Implementation

### 2.1 Data Flow



# 2.2 PostgreSQL/PostGIS

PostgreSQL is an open-source object-relational database management system (DBMS) known for its robustness, scalability, and extensive features. It is often referred to as "Postgres" and is one of the most popular and widely used databases in the world. PostgreSQL supports a wide range of data types, including numeric, text, Boolean, date/time, JSON, XML, and more. It provides support for complex queries, indexing, and advanced features such as full-text search, spatial data storage and querying, and transactional processing with ACID (Atomicity, Consistency, Isolation, Durability) properties.

PostgreSQL follows the SQL (Structured Query Language) standard, and it also provides additional features beyond the standard SQL specification. It supports stored procedures, triggers, and views, allowing developers to define custom business logic within the database itself.

PgAdmin is an open-source administration and development platform for PostgreSQL. It is a graphical user interface (GUI) tool that provides a convenient way to manage and interact with PostgreSQL databases.

**PostGIS:** PostGIS is an open-source spatial database extension for PostgreSQL. It adds support for geographic objects and spatial functions to the PostgreSQL database, enabling the storage, retrieval, and analysis of geospatial data. Geographic data such as points, lines, polygons, and multi-dimensional geometries can be stored and manipulated within your PostgreSQL database using PostGIS. The capabilities of the database are extended by PostGIS to handle spatial data types, indexing, and spatial operations. PostGIS has gained popularity in a variety of applications, including mapping, geolocation-based services, environmental analysis, urban planning, and transportation. Its combination with the power and versatility of PostgreSQL makes it a robust solution for managing and analyzing geospatial data in a relational database environment.

The below image shows the PGadmin tool.

irowan 🗴 🎹 🐔 Q	Dathboard Properties IICA, Italictics	Dependencies Dependents	
· El Servera (5)	1.4		
+ ELKEPORK	Tibe	TYLETTIE	Reightagon
<ul> <li>cf Kal-park project MAETER</li> </ul>	- This	RETURN OV size retrain prove	normal
<ul> <li>ET Kat-pain project READER</li> </ul>			
• If PodgoodCL 13			
- Q style			
<ul> <li>Brownows IIO</li> </ul>			
- Revis			
> @Cents			
> -9/ Caterous			
<ul> <li>Event Triggers</li> </ul>			
> 16 Extensions			
Foreign Data Wramerk			
+ //JLanguages			
- Wilchernag (7)			
<ul> <li>B-DH</li> </ul>			
<ul> <li>Improvement</li> </ul>			
<ul> <li>epiblic</li> </ul>			
~ ÷ 100			
a) di Constione			
Hg Dornains			
<ul> <li>IB FT6 Configurations</li> </ul>			
<ul> <li>IN ITS Dictionaries</li> </ul>			
<ul> <li>Au FTS Parsers</li> </ul>			
<ul> <li>IEFTS Tempateo</li> </ul>			
<ul> <li>IEPPorsign Tables</li> </ul>			
<ul> <li>MiFunctions</li> </ul>			
MaterialIed Views			
* Theorem and			
> 1.1Sequences			
<ul> <li>Raties (6)</li> </ul>			
+ Ittige, Matters			
+ ministration			
+ Hitkatalog_stoff			
<ul> <li>Immessionemetritit</li> </ul>			
<ul> <li>Immensatele</li> </ul>			
<ul> <li>Eliminativent</li> </ul>			
<ul> <li>W Tripper Functions</li> </ul>			
> 10 Tupes			
- 10 Yiews (7)			
<ul> <li>Boopty_station_result</li> </ul>			
e 📷 nitat			
+ iteaLgeon			
		image 2- PGadmin	

Watching the below videos to understand how we can create schemas and tables in the PostgreSQL database.



Create a database and schema based on the above video.

# 2.3 Data Engineering

# 2.3.1 Downloading the Data

In the first step, The data must be downloaded from here. To complete the process, download and extract the initial zip file, which contains four CSV files and one set of instructions. Refer to image 3 for guidance on selecting the appropriate zip file.

		information und Hondrhain West	Technik Mar
OpenGeodata.NRW J Dryges:		(1 <b>3</b>	ion (* XML
Grundwassermessstellen NRW			
Linesteinformetion: Diese Dalen siehen unter der Dekentitanz Deutschland - Zero - Veral     Delensatzbeschreibung: [2] Operingshitz, meta Zep     Weitadeten im [2] Opering 2010	en 2.8. Jede Nutzung ist ohr	e Shechtikkungen nder Bedingunge	et dallang
Deleter des Onie Responseries establice la MORIFICIA (ORIFI)			
Daten der Grundwassermessstellen in NRW als GSV			
Datien der Grundwassermessstellen in NRW als CSV		Letzte Änderung	Dataigroße
Datel Detel		Letzte Änderung 25-10.2021 09-48-00	Date-igrofie 41.7 MB
Date: Der Grundwassermessstellen in NKW als CSV Date:  © OpentigsteG, gw-messdellen-messwerte, EPS625002, CBV sp  © OpentigsteG, gw-weserstand, EPS625002, CBV sp		Letzte Änderung 28-10 2021 08-46.00 29-10 2021 08-49-00	Datelgrolle 41.7 MB 100.4 MB
Daten der Grundwassermessstellen in NKW als CSV Detail  DentfligteC, gwmeastellen-messweis_EPSG2802, CSV zp  (d) OpenfligteC, gwweeserstel, EPSG2802, CSV zp Daten der Grundwassermessstellen in NRW als Sqilte Datenbank		Letzte Änderung 29-10.2021 08-48-00 29-10.2021 08-49-08	Dataigrófia 41.7 MB 100.4 MB
Daten der Grundwassermessstellen in NRW als CSV Detai  Deten der Grundwassermessstellen in NRW als Sqilte Datenbank Detai		Letzte Änderung 28-10.2021 08-48.00 29-10.2021 08-49:08 Letzte Änderung	Dataigróile 41.7 MB 100.4 MB Dataigróile
Daten der Grundwassermessstellen in NKW als CSV  Date		Letts Anderung 28-10.2021 08-48.00 29-10.2021 08-49:00 Letzts Anderung 29-10.2021 08-49:25	Dateigröße 41.7 MB 300.4 MB Deteigröße 88.6 MB
Daten der Grundwassermessstellen in NKW als CSV  Datel  Deten der Grundwassermessstellen in NRW als Sqlite Datenbank  Datel  Opentrigstof, ge-messstellen-messvenz, EPS02602, Sqlite zp  Dotel  Detei  Deteil  Deteil		Letts Anderung 28-10.2021 08-48.00 29-10.2021 08-49:00 Letzts Anderung 28-10.2021 08-49:25 29-10.2021 08-49:25	Dataigrolle 41.7 MB 100.4 MB Dataigrolle 88.8 MB 205.1 MB
Daten der Grundwassermessstellen in NKW als CSV  Detei  C OpentrigsteC, gwinnesstellen messwerte, EPSG2802, CSV zip  Daten der Grundwassermessstellen in NRW als Sqlife Datenbank  Datei  CopentrigsteC, gwinnesstellen inserwerte, EPSG2882, Sqlite, zip  CopentrigsteC, gwinnesstellen in SRW als Shape		Letzte Änderung 28-10.2021 08-48.00 29-10.2021 08-49-08 Letzte Anderung 28.10.2021 08-49-20 28-10.2021 08-50:11	DateSprote 41.7 MB 100.4 MB DateSprote 86.9 MB 206.1 MB
Daten der Grundwassermessstellen in NKW als CSV  Detei  C OpentrigsteC, gwinnesstellen in NKW als Sqlife Detenbank  Detei  OpentrigsteC, gwinnessstellen in NKW als Sqlife Detenbank  Detei  OpentrigsteC, gwinnessstellen in SpSG2882, Sqlite zp  Grundwasserabhängige LandSkosysteme in NRW als Shape  Detei		Letzte Änderung 28-10.2021 08-48.00 29-10.2021 08-49-08 Letzte Änderung 28-10.2021 08-49-08 28-10.2021 08-49-29 28-10.2021 08-5014 Letzte Änderung	Detelgrothe 41.7 MB 100.4 MB Detelgrothe 86.9 MB 306.1 MB Detelgrothe

After the above zip file has been extracted, the four CSV files and the instruction file can be considered, with a strong recommendation to read the instruction file first.

# 2.3.2 Python

Python is a high-level, interpreted, and general-purpose programming language known for its simplicity and readability. It was created by Guido van Rossum and first released in 1991. Python emphasizes code readability and has a design philosophy that emphasizes clear, concise syntax, making it easier to write and understand code. Python is used in several ways such as:

- AI and machine learning
- Data analytics
- Data visualisation
- Programming applications

In this project, we have used Python for data engineering, data pre-processing, and data analysis. Jupyter Notebook is used in this project to write Python codes. The Jupyter Notebook is an opensource web application that data scientists can simply write the code for and make it easier to document. Simply, we can combine Python codes, text, images, comments, and the result of the codes on the same page. The below image shows how code, text, and the result of the code can be seen on a single page.

Eiles E	-	in in	nert Call	Warnel A	distante.	Links						Texateri	Dethon 3 O		
- 100 L			Gen Cen	Contrast 1	magaze	1000						1.0000000	Trymura O		
四 +	9< P)	15 +	🔶 🕨 Run	E C »	Markdow	ei Y	523								
	F	Read t	he file		>	Text									
	0	on't forge	t to add the "de	elimiter", nom	nal csv file	's delimiter i	is "," but in ou	ur case it's *;*	, so we need	to assign it.					
	_							2222222222							
	100								_						
In	[59]: d	f = pd.r f.bead()	ead_csv(gw_s	tation_pfna	ame, deli	imiter=';'	', header='	infer')			Python Code				
In	[59]: d	f = pd.r f.head()	read_csv(gw_s	tation_pfna	ame, deli	imiter=';'	', header='	infer')			Python Code				
In   Out	59]: d	f = pd.r f.head() sl_m	read_csv(gw_s	tation_pfna	ane, deli •32	imiter=';' n32 gw	', header='	infer') grundstueck	gemeinde_id	gwhorizont_id	Python Code gwhorizont	beobachtung	_wasserstand	-	
In   Out	[59]: d	f = pd.r f.head() sl_m 0 67530	read_csv(gw_s messstelle_id 32505929	tation_pfna name UWB-Ddorf 01285	•32 343064 1	imiter=';' n32 gw 9578019	", header=" w_stockwerk 1.0	infer') grundstueck	gemeinde_id 05111000	gwhorizont_id NaN	Python Code gwhorizont NaN	beobachtung	_wasserstand	٦.	
In   Out]	[59]: d d	f = pd.r f.head() sl_mr 0 67530 1 51044	read_csv(gw_s messstelle_id 32505929 10446746	name UWB-Ddorf 01285 60GP012303	•32 •32 343064 ! 292077 !	miter=';' n32 gw 9678019 9645349	", header=" w_stockwerk 1.0 NaN	infer') grundstueck privat	gemeinde_id 05111900 NL000882	gwhorizont_id NaN	Python Code gwhorizont NaN Zwischermittel	beobachtung	_wasserstand	₽	
In   Out	59]: d d	f = pd.r f.head() sl_m 0 67530 1 51044 2 51070	read_csv(gw_s messstelle_jid 32505929 10448746 87005323	NUMB-Delart UVWB-Delart 01285 600F012303 588F024600	•32 •32 343064 1 292077 1 287141 1	m <mark>32 gw</mark> 5578019 55845349 5584535	", header=" w_stockwerk 1.0 NaN NaN	infer') grundsturck privat	gemeinde_id 05111000 NL000882 NL001640	gwhorizont, id NaN 5 00	Python Code gwhorizont NaN Zwischermittel Neurather Sand	beobachtung	wassenstand	₽	

### 2.3.3 Anaconda

Anaconda is an open-source distribution for python and R. It is used for data science, machine learning, deep learning, etc. With the availability of more than 300 libraries for data science, it becomes fairly optimal for any programmer to work on anaconda for data science Anaconda is used in this project. An environment on Anaconda has been created to install all the packages needed for this project.



**Environment in Anaconda:** A conda environment is a directory that contains a specific collection of conda packages that are used in the project.

**How to create Conda environment:** The below video shows how to create a Conda environment, how to activate it, how to install different packages on the environment and how to deactivate the environment.



**Python packages:** Python packages are collections of modules that provide additional functionality and tools to extend the capabilities of the Python programming language. Packages are typically distributed and installed using package managers such as pip (the default package manager for Python) or conda.

To get more details about conda environment, I highly recommend visiting the below webpage.

https://towardsdatascience.com/manage-your-python-virtual-environment-with-conda-a0d2934d5195

**openhype environment:** For this project, an "openhype" environment was created to install all the required packages. This environment ensures that all the necessary dependencies are properly installed and configured.

Several packages are related to data science but in this project, we have used the below packages. There are two ways to install the below packages:

**install a package manually:** In this way you need to install each package manually into openhype environment by the command prompt.

• **pandas:** Pandas is a popular open-source Python library for data manipulation and analysis. It provides easy-to-use data structures, such as DataFrame, Series, and Index, that are designed to handle structured data efficiently.

In this project, pandas was utilized to read the CSV files, clean the data, and perform data engineering tasks. To install Pandas, execute the following Python code within the openhype environment in the Anaconda prompt:

conda install pandas

• **sqlalchemy:** SQLAlchemy is a popular open-source SQL toolkit and Object-Relational Mapping (ORM) library for Python. It provides a set of high-level APIs that allow developers to interact with relational databases using Python code.

conda install sqlalchemy

• **psycopg2:** Psycopg2 is a PostgreSQL adapter for Python. It provides a Python interface for interacting with PostgreSQL databases, allowing developers to connect to a PostgreSQL database server, execute SQL queries, and perform database operations using Python code.

conda install psycopg2

• geopandas: Geopandas is an open-source Python library built on top of Pandas and other

geospatial libraries. It extends the capabilities of Pandas by adding support for geospatial data, enabling users to work with spatial data in a tabular format.

conda install --channel conda-forge geopandas

Certain packages require the specification of a channel for installation. This is why the above code includes the channel specification to ensure the correct installation.

• **jupyter notebook:** Jupyter Notebook is an open-source web-based interactive computing environment that allows users to create and share documents containing live code, visualizations, explanatory text, and more. It supports various programming languages, including Python, R, and Julia.

conda install jupyter notebook

To ensure the successful installation of all the aforementioned packages, it is crucial to install them within the openhype environment using the Anaconda prompt.

**Load all the packages into the environment by a YAML file:** To streamline the installation process for all the required packages in this project, it is recommended to create an environment named "Openhype" and load a YAML file containing the package specifications. The contents of the "openhype.yml" file, as shown in the image below, encompass all the necessary packages.



The following code allows you to create an environment based on the yml file. Keep in mind that the YAML file should be in the same directory as your Anaconda installation. If the file is located elsewhere, you will need to provide the full path to the file in the code. openhype.yml can be found from here

```
conda env create -f openhype.yml
```

How to create a yml file for the environment: With the below code you can export the yml file from the existing environment.

conda env export > openhype.yml

Since the four CSV files have been downloaded in the previous chapter, it is time to read the CSV files and initiate the cleaning process to make them ready for importation into our database.

Refer to the code section in our Notebook for more information Github (the below image).

Search or jump to	Pull requests Issues Marketplac	e Explore			0 +• <b>6</b> •
sina7272/LANUV_Groundwater	(Refe)			Schin ⊕ Unweich I + 17 ( )	🚼 Shaved (1) 🔹
O Code 💿 Issuer 💠 Poll requests	🔆 Actions 🗄 Projects 💷	Wild 😳 Security 🖂 Insights 🔅	Settings		
	Prese Pterest 00		Go to file Add file - Code -	About Ø	
	ates 7777 and ate		Antibility of James and 10 22 another	Ground water in NRW	
			and the Gamma		
	Apyrib, checkpoints			Q total Q teatering	
	Natebook				
	b data		17 days ago		
	· 978			Releases	
	D glattidutes				
	D README.md				
	README.md			Packages	
				No and American American	
	LANUV_Groun	dwater			
	Ground water in NRW			Languages	
				Deside Relation 12.70 P. 406.025	
1					

There below four notebooks should be run separately, in order to import data into the database.

• import\_gemeinde.ipynb:

In this notebook, we will import the data of all geminde into the database.

• import\_katalog\_stoff.ipynb:

In this notebook, we will import the data of all the catalogue substances into the database.

• import\_messstelle.ipynb:

In this notebook, we will import the data of all stations into the database.

• import\_messwert.ipynb:

In this notebook, we will import the data of all values into the database.

# 2.4 Observation Data in the Database

After successfully downloading, cleaning, and importing the data into the database in the previous section, it is now time to examine the data within the database. Since we are working with a PostgreSQL database, we can utilize SQL commands to retrieve and analyze the data. Let's run some basic SQL commands to gain insights into the data.

To view our tables, we can use the following code. It selects all the columns from the "hygrisc" schema, which contains our tables. We limit the output to the first 100 rows to avoid overwhelming results.

select \* from hygrisc.messwert limit 100;

Image 7 illustrates the outcome of the aforementioned command executed in PGAdmin. It displays the resulting data obtained from the execution of the provided SQL command.

	5 III 1 Q	Dashbo	and 1	Properti	es SQL	Statis	tics De	ependenc	ies i	Dependents	6	etwais/etwa	is@alw	105 *												
	> lig Domains	8 8		× 8	E Q.		~ 5	8 0	8-	Τ -	N01	init 🖂			4.0	- 1	5	6	dr'	£	<b>8</b> -					
	<ul> <li>ETS Configurations</li> </ul>	d	was/sh	vooSelv	A36 -	ć.																				
	» DA FTS Dictionaries	Ouers F	other.	Duary I	lictory																	Scratch i	Port			
	Aa FTS Parters     To Toroniston	term y a			tanti j																	-renditer i	-			
	> Francisco Tables	1.5	elect	+ from	s sina.re	essaer	t Limit	t 100;																		
	bitFunctions																									
	> Materialized Views																									
	> (() Procedures																									
	<ul> <li>L1 Sequences</li> </ul>																									
	<ul> <li>Tables (6)</li> </ul>																									
	> mgw_stations																									
	> ⊞katalog_ge																									
	> 🔠 katalog_stoff																									
	measurements																									
	> 📑 messstelle																									
	> et messert																									
	messistelle     messivert     Digger Functions	Data Or	tout /	Explain	Messar	os N	officatio	08																		
	S In messatelie     S In messatelie     S Integer Functions     Types     Types	Data Os	aput	Explain	Messag	es N	lotificatio	n))		100																
	If messatelie     If messatelie     If messateri     Winger Functions     Types     Week (7)	Data Os	itput ILer PKI bior	Explain	Messag messstelle_i bont	es N	otificatio	na 1 data	npit	stoff,/w	/	proberigut	1	nsergebr	tilla_m talion	1	nesserg	pitric.)	inveis	,	bestimmur double pre	gagrenze	,	massesthett	,	trerewerfahren brot
	(messatelie     (if messatelie	Data Os	itput il.er (PK)biger 12	Explain	Messag messstelk_j bigint 295	es N d /	iotificatio pra_id Not 3/1092/1	na data data 16. 1992	n.jan 05-21	≠ stoff,.w bight	1	probergut text	1 10	nsengebr uble preci	iston 105	/ 1 243 II	nesserg Art	peteria_1	invelis	,	bestimmu double pre	igagienze clatori	/	massesthet text mg/l	,	tnereworfahren text Meniscanfiber
	> imessatele     > imassert     > ignager Functions     > if Types     v     Versi(7)     > ignosplo_station_result     > initiat     v     initiat.secm	Data Or	etput et.or (PK)bigin 12 12	Explain 4 299051 299053	Messag messstelle, j tagint 295 200	es N d /	preLid Not 3/1992/1	na datar data 16. 1992	<b>n., pin</b> -05-21 -03-29	≠ stoff.w bight	1244	probersput text Grundwatser Grundwatser	1 10	nssergebr uble pred	ia_e teron 105: d1/	248 ji	nesserg Art Full)	pitria_t	inveis	,	bestimmu double pre	ngsgrente chilon	1	massembelt text mg/l mg/l	,	transverfahren text Membranfiher Membranfiher
	imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     imess(7)     imess(7)	Data Or	itput iLer PK]bigin 12 12 12	Explain 4 209051 209053 209053	Messag messstele.j tagint 299 200 80	es N d /	otificatio pre_id Not 3/1992/1 3/1994/1 3/1993/1	na datur date 16. 1992 16. 1994	n.pn -05-21 -03-29 -01-11	/ stoff.w	/ 1244 1244	probesgut teat Grundwasser Grundwasser	/ m	nssergebr uble preci	Haus Islan 105: 61/ 3.0	248 ji 978 ji 999 ji	nesserg art rull) rull)	pitria_1	inetis	,	bestimmu double pré	ngagi vertate ctalizati	int int	massembelt text mg/l mg/l mg/l	,	trensverfahren toot Menipranfiker Menipranfiker Nerstoranfiker
	<ul> <li>immessatelie</li> <li>immessent</li> <li>iterations</li> <li>Types</li> <li>Weens (7)</li> <li>iteration (7)</li> <li>i</li></ul>	Data Os	djor PKI bigen 12 12 12	Explain 4 209051 209053 209053 209072 200077	Messag messatele, i lagint 299 209 209 80 80 80 80 80 80 80 80 80 80 80 80 80	es N d / 1002710 1002916 1900097	otificatio prauid text 3/1992/1 3/1993/1 3/1993/1	na datar data 16. 1992 16. 1993 16. 1993	05-21 -03-29 -01-11 -11-02	/ stoff.nr bight	/ 1244 1244 1244	probesgut text Grundwasser Grundwasser Grundwasser	1 10	nsergebr	Mae Iston 104: 3.0 2.25	<ul> <li>* 1</li> <li>248 11</li> <li>978 11</li> <li>978 11</li> <li>979 11</li> </ul>	nesserg Art Full) Full) Full	ebris.	inetit	,	bestimmu double pre	ngagnentes ctation	Find Study Study Study	masseathait teat mg/l mg/l mg/l mg/l	,	transvarfahran text Membranfilter Membranfilter Membranfilter
	messatele     messatel     messatel     messatet     messatet     groups_station_result     mess(7)     more groups_station_result     minat     montat_peen     plates     messatele     messatele     messatele     messatele	Data Or 1 2 3 4	dput eLer PK(bige 12 12 12 12 12	Explain 4 209051 209053 209072 209077 209077	Messag messstele.i tagint 299 200 80 80 80 80 80 80 80 80 80 80 80 80 8	es N d / 1002710 1002916 19002916 1900303	Intificatio peal.id Net 3/1992/1 3/1993/1 3/1993/1 3/1992/1 3/1992/1	na datar date 16. 1992 16. 1993 16. 1993	-05-21 -05-21 -03-29 -01-11 -11-02 -10-22	<ul> <li>stofLar</li> <li>bipint</li> </ul>	/ 1244 1244 1244 1244	probesigat Seat Grandwasser Grandwasser Grandwasser Grandwasser	/ ma do	nseigebi uble preci	Haue 106: 01/ 3.0 2.25 3.05	243 (1 978 (1 978 (1 977 (1) 977 (1) 903 (1)	nessen set cull cull cull cull	patienia_1	inville	,	bestimmu double pre	ngagmentas chilitan	<ul> <li>Internet</li> <li>Internet</li> <li>Internet</li> <li>Internet</li> <li>Internet</li> <li>Internet</li> </ul>	masseathait text mg/l mg/l mg/l mg/l	,	transvarfahran toot Menipranfilter Menipranfilter Menipranfilter Menipranfilter
	imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     impers     imperseteeeeeeeeeeeeeeeeeeeeeeeeeeeeeeeeee	Deta Or 1 2 3 4 5	2001 6Lor (PK) biger 12 12 12 12 12 12 12	Explain 4 209051 209053 209072 209077 209079	Messag messstelle, i tagint 20% 80 80 80 80 80 80 80 80 80 80 80 80 80	es N 602710 6022710 6022916 1302097 1302093	otificatio peal.id fact 3/1992/1 3/1993/1 3/1993/1 3/1993/1	na data 5ate 16. 1992 16. 1993 16. 1993 16. 1992	n.pn 05-21 03-29 01-11 11-02 -12-22	<ul> <li>stoff_re</li> <li>bigint</li> </ul>	/ 1244 1244 1244 1244	probesgut Seat Grundwasser Grundwasser Grundwasser Grundwasser		niseigebr	104. 104. 104. 3.0 2.25 3.05	243  1 243  1 978  1 978  1 989  1 777  1 403  1 625  1	1455007 Art Gall Gall Gall Gall Gall Gall	pitris.)	invels	,	bestimmu double pre	ngagmentas chaiten	Start Start	masseshielt text mg/l mg/l mg/l mg/l mg/l	,	transvarfahren text Membranfiher Membranfiher Membranfiher Membranfiher
	imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     initiat_pectn     initiat_pectn     initiat_pectn     initiat_pectn     imessatelie     initiat_pectn     imessatelie     imessatelie     imessatelie     imessatelie     imessatelie     imessatelie	Deta Or 4 3 4 5 6	2004 2007 22 12 12 12 12 12 12 12 12 12 12 12 12	Explain 1299051 299053 299072 299077 299079 299081	Measag messatela.i tagin 20% 80 80 80 80 80 80 80 80 80 80 80 80 80	es N d / 002710 002710 13002710 13002710 1300303 1300303 1300303	etificatio preid text 3/1992/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1	na data data 10. 1992 10. 1993 10. 1993 10. 1993 10. 1993 10. 1993 10. 1993 10. 1993	05-21 03-29 01-11 -11-02 -12-22 04-21 -03-10	/ stoff.av bigen	/ 1244 1244 1244 1244 1244	probesgut Seat Grundwasser Grundwasser Grundwasser Grundwasser Grundwasser	1	nsergebr	108.2 108. 108. 108. 11.0 11.0	248 11 978 11 978 11 977 11 9039 11 9039 11 9039 11 9039 11	nesserg ext cull cull cull cull cull	pitric.)	inveis	,	Sestimmu double pre	ngagrenta ctalion		massembelt text mg/l mg/l mg/l mg/l mg/l mg/l mg/l	,	transvarfahren text Mentbranfiker Mentbranfiker Mentbranfiker Mentbranfiker Mentbranfiker
	<ul> <li>messatele</li> <li>messatel</li> <li>messatet</li> <li>messatet</li></ul>	Deta Ou 1 2 3 4 5 6 7	tput sl.or PK[bigit 12 12 12 12 12 12 12 12 12 12	Explain 1299051 299053 299072 299077 299079 299081 299097	Messag messetele, i tagin 20% 80 80 80 80 80 80 80 80 80 80 80 80 80	es N d 002710 002710 002710 1002700 1002700 10000000 10000000 10000000 10000000 10000000 10000000 100000000	totificatio pealid text 3/1992/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1	<ul> <li>datus date</li> <li>10. 1992</li> <li>10. 1993</li> <li>10. 1993</li> <li>10. 1993</li> <li>10. 1992</li> <li>10. 1993</li> <li>10. 1993</li> </ul>	05-21 03-29 01-11 11-02 12-22 04-21 07-19	/ stoff_m bight	/ 1244 1244 1244 1244 1244	probesgut text Grandwasser Grandwasser Grandwasser Grandwasser Grandwasser Grandwasser		essergebr	108.0 108. 3.0 2.25 3.05 11.0 106.	243 11 243 11 978 10 978 10 977 10 603 10 675 10 248 10	nesserg ext cull cull cull cull cull cull	petimic.3	invels	,	bestimmur double pre	gsgrente claton		messenheit tert mg/l mg/l mg/l mg/l mg/l mg/l	,	trensverfahren bot Menistanfäher Menistanfäher Menistanfäher Menistanfäher Menistanfäher Menistanfäher
	<ul> <li>immessatelie</li> <li>immes</li></ul>	Deta Ou 4 1 2 3 4 5 6 7 8	tput sl.w PK bige 12 12 12 12 12 12 12 12 12 12	Explain 209051 209053 289072 289077 289079 289081 289097 289099	Messag messetele, i tagin 20% 80 80 80 80 80 80 80 80 80 80 80 80 80	es N d 1002710 1002703 1002703 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 10027515 1002755 1002755 1002755 1000000 1000000 10000000 10000000 10000000 10000000 10000000 100000000	otificatio pealid text 3/1992/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1 3/1992/1	<ul> <li>datus date</li> <li>10. 1992-</li> <li>10. 1993-</li> </ul>	05-21 -05-21 -03-29 -01-11 -11-02 -12-22 -04-21 -07-19 -05-10	stoff.or	/ 1244 1244 1244 1244 1244 1244	probesput text Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser		essergebr	104. 104. 104. 104. 104. 104. 105. 11.0 105. 53.	243    243    978    989    989    903    675    248    124	nesserg éxt cull cull cull cull cull cull cull cul	pinis,	itterellik	,	bestimmur double pre	gsgrente claton		massemheit text mg/l mg/l mg/l mg/l mg/l mg/l	,	tereverfahren tot Mentschliter Mentschliter Mentschliter Mentschliter Mentschliter Mentschliter Mentschliter
,	<ul> <li>immessatelie</li> <li>immes</li></ul>	Deta Ou 1 1 2 3 4 5 6 7 8 9	rput eLev (PK)bigen 12: 12: 12: 12: 12: 12: 12: 12: 12: 12:	Explain 4 209051 209055 209052 209077 209079 209081 209097 209090 209009 209105	Messag messtela_ 291 201 80 80 80 80 80 80 80 80 80 80 80 80 80	es N 002710 002710 002916 1300303 1300303 1300303 1300364 1300364 1300364	otificatio pha.id 14xt 3/1992/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1 3/1993/1 3/1992/1 3/1092/1 3/1002/1	rtal datar dat	05-21 03-29 01-11 -13-02 04-21 07-10 06-10 07-14	stoff.ar	/ 1244 1244 1244 1244 1244 1244	probesput text Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser Dranderasser		nsergebi	his_e 1900 01/ 3.0 2.25 3.05 11.0 106 53 9.7	248    248    989    777    803    875    843    124    124    124	1455000 601 601 601 601 601 601 601 601 601	Coindag	inweis V Su	/	Sestimanus double pre	gagrenze claion	Part Part Part Part Part Part Part Part	massemheit text mg/l mg/l mg/l mg/l mg/l mg/l mg/l mg/l	,	terrverfahrer tot Mentschlier Mentschlier Mentschlie Mentschlie Mentschlier Mentschlier Mentschlier Mentschlier

image 7- messwert table in databse

Now with the above SQL command, we are able to see the other three tables that we have (The below codes).

select \* from hygrisc.messstelle limit 100;

select \* from hygrisc.katalog\_stoff limit 100;

select \* from hygrisc.katalog\_ge limit 100;

Now we want to see more details for our tables and we will run the below codes.

**Count the rows of each table:** With the below code, we can see how many rows we have in each table.

select count (\*) from hygrisc.messwert;

select count (\*) from hygrisc.messstelle;

**Filter the data based on Nitrate only:** To determine the substance number of Nitrate, we need to retrieve the corresponding information from the database.

select \* from hygrisc.katalog\_stoff where name like 'Ni%';

<ul> <li>w period</li> </ul>	8 8 8 - E Q - 8 - 5 E Z- T - mm - E	5 5 4 A R-
- (b Stra		
a di fontano		
1 Diffe Continuations	Coarty Entern Owey Hillingy	SCHED PHE.
+ In FTS Dictionwies	1 select + from size Astalog_stoff where name 15kg 'Nrth'	
<ul> <li>Au FTS Pachers</li> </ul>		
+ III FTS Templates		
<ul> <li>Ittl: Foreign Tables</li> </ul>		
> Bit Punctions		
<ul> <li>Materialized Value</li> </ul>		
<ul> <li>#"pProcedures</li> </ul>		
+ 1.1 Sequences		
- Tables (6)		
<ul> <li>Itigx_stations</li> </ul>		
<ul> <li>Ethelwing.get</li> </ul>		
> mikataing_stuff		
<ul> <li>III measuramanta</li> </ul>		
* mirrennitelle		
<ul> <li>III messavet</li> </ul>		
<ul> <li>BQ Tripper Functions</li> </ul>		
* Dillibert		
a second state with the count	Dete Output Explain Messages Nothcattone	
a contract of the second second	statur a tunte a beat-stature a celul a	
- Prittat asom	a bigm - wert - double precision - weit -	
+ D Columna	1 1143 Non pull 746-83-1	
+ ma Robert	2 THE NOME INT THE CO	
+ 2+ triggers	8 1244 Miner Evall Sould	
+ mattat	<ul> <li>4 1240 Weindtelandf.</li> <li>[54] [54]</li> </ul>	
+ safat_peom	5 T246 Meyer (mill 2mill	
+ 🗰 terrar	6 1047 tem Stolentt (rull, jrull)	
+ 🚺 terni2	7 197 Netternite Tessie (rull, trail,	
F 🐵 tiger	8 2293 Nexteenal (will 19-03	
<ul> <li>Inper_data</li> </ul>	b 2042 Months [out 129-365]	
> (e) fogology	10 2457 Internetworke+15%	
<ul> <li>m pestiples</li> </ul>	11 THE MORE THE STREET	
<ul> <li>Eternoliata_postors</li> </ul>	41 Martin Martine Andreas	
<ul> <li>Therman</li> </ul>	The state of the s	
1. The second diversion diversion	12 THE RELEASE (10) 20(5)	
<ul> <li>The Login / Broup Holes</li> </ul>	No. 1 August and a second s	

According to the information presented in "Image 8," the substance number (stoff\_nr) of Nitrate is **1244** 

select \* from hygrisc.messwert where stoff\_nr = '1244';

Now we can filter the messwert table based on Nitrate. In this step, We can save this new table as a new "view".

**What is a view:** a view is a virtual table that is derived from one or more existing tables or other views. A view does not store data physically but rather provides a way to present data from underlying tables in a structured and organized manner. It acts as a predefined query that can be used to retrieve and manipulate data. The below code creates views:

```
create view hygrisc.nitrat as (select * from hygrisc.messwert where stoff_nr
= '1244');
```

We now have a "nitrat" view, which can be accessed just like a table using the following code. This view is filter of our messwert table based on "1244" which is "Nitrate"

select \* from hygrisc.nitrat ;

**Group by the two tables:** the Group by clause is used to group rows based on one or more columns in a table. When working with two tables, you can perform a GROUP BY operation to group the data based on common values from both tables. In this section, we want to group by our two tables (messwert and messstelle) only in Nitrate. These two tables have a column **messstelle\_id** which means station id.

```
select messstelle_id, count(*) from hygrisc.messwert where stoff_nr =
'1244' group by messstelle_id;
```

a 10 public		
- (0.10a		A B
+ BL Collefierza	g steas/daasgetass -	
+ 4 Domana	Overv Diltor Overv History	Scratch Ped
+ D.FTS Configurations	The second	
+ (N. F75 Dictionates	I Select mesostelle_id, count(+) from sine.mesowert where stoff_nr + '1244' group by mesostelle_id	
<ul> <li>Av FTS Parsers.</li> </ul>		
+ 🛄 FTS Templates		
<ul> <li>W Foreign Tables</li> </ul>		
+ No functions		
+ 😻 Materialized: Views		
<ul> <li>Upprocedures</li> </ul>		
<ul> <li>* 1.3 Seduences</li> </ul>		
- Ell Tatsies (6)		
<ul> <li>Illigik_stations</li> </ul>		
s mikabalog.ge		
> micatalog_stoff		
<ul> <li>Esmeasurements</li> </ul>		
<ul> <li>Internationality</li> </ul>		
s III metawen		
<ul> <li>Big trigger rundhard</li> </ul>		
- Marriell		
a consider station most	Della Output: Explain: Neosages: Notifications	
s contrast	recepted a rout a	
- Control peope	A phu _ phu _	
+ In Columna	1 10151210 34	
e em Fisien	2 123040 3	
+ 2+ Tragers	E 18290411 28	
* Castat	4 1020001 28	
<ul> <li>multid_peore.</li> </ul>	8 162842716 IN	
> Tomp	6 1020000 20	
s 🔚 terral	T 100040 J	
<ul> <li>Hotiger</li> </ul>	a 100100 at	
<ul> <li>Wittger_data</li> </ul>	4 102117 47	
> (e) topology	N NATION OF	
> 🖾 boepbeer	10 10 10 10 10 10 10 10 10 10 10 10 10 1	
<ul> <li>Brandela, postars</li> </ul>	1 UNDER 00 00	
> m tearap	16 Institute at	
<ul> <li>Todau Australia Hoseit</li> </ul>	13 1621312 4	<ul> <li>Boccessfully run. Total query runtime: 471 maer. 2632 room affectail.</li> </ul>
<ul> <li>Tatoxipacwo</li> </ul>	E 14 11201340 12	

Image 9 shows that each station id has how many single measurements for the Nitrate only. I highly recommend opening the below website to get more deep into how "group by" works and how we can use it.

https://www.w3schools.com/sql/sql\_orderby.asp



#### Maximum date in nitrat table:

select \* from hygrisc.nitrat where datum\_pn = (select max(datum\_pn) from hygrisc.nitrat); 2025/01/19 14:49

rowser S III %	Q, Dathboard Properties 30, Statistics Dependencies Dependences and insurvivous/stress."		×
• • • • • • • •	8 8 8 - 2 9 - 5 - 5 E 2 - T - 100 - 8 8 - 5 8 - 5 8 - 5 8 - 5 8 -		
> \$1 constores	6 Johnas/shreesgotnes +		
+ Rp Domaini	Guary Editor Guary History	Scratch Pad	
Q FTB Configurations     Virg TS Configurations     A TS Parami     A TS Parami     Fill Tangatotic     Virg Tangatotic     Virg Tangatotic     Fill Tangatotic     F	I select + from sins.mitrat where detum_on + (select max(detum_on) from sins.mitrat)	2018/01/198	
> 10 Toget Functions > 0 Vrest - 0 Vrest - 0 0routper, station, result + 0 router + 0 0routper, station, result + 0 router + 0 rou	Cetto Organ         Definition         Massach         Massach	writers o scort o refuel to as     socialize interview as     socialize interview as     socialize interview as	•

As we can see in Image 10, the maximum date is 2021-08-17

#### Minimum date in nitrat table:

select \* from hygrisc.nitrat where datum\_pn = (select min(datum\_pn) from hygrisc.nitrat);

a and a second s	🎹 🐂 🔍 Dashboard Properties 203. Etablicitz Dependencies Dependencies 👩 elway/elwast/		8
<ul> <li>se pated</li> </ul>	S & B Y E D Y B Y E B BY T Y THE Y E B Y A D Y S S AY A BY		
·			
+ S.Conatione	Construction of the second sec		
<ul> <li>BUTT Contractions</li> </ul>	Overy failor Overy failors	Soutch Pad .	,
<ul> <li>M FTS Dictionaries</li> </ul>	3 select + from wine, mitrat where dotum_pn = (select minidatum_pn) from wine, mitrat);		
+ #4 FTS Paraelo			
<ul> <li>FTS Templates</li> </ul>			
> III Foreign Tables			
<ul> <li>BiFunctions</li> </ul>			
<ul> <li>Other Mathematical Views</li> </ul>			
> \$1Procedures			
<ul> <li>1.3.Sequences</li> </ul>			
- CTabres (S)			
<ul> <li>Engles_dations</li> </ul>			
<ul> <li>Imitiation shall</li> </ul>			
<ul> <li>Immenostate</li> </ul>			
+ III measured			
> 89 Trigger Functione			
> C Types			
- 📫 Viens (7)	Data Output Explain Messages Notifications		
+ proupto_atation_result	al ar massada id una id datem ar and ar protector massadates o tassadates lineas bettern recovers massadate treventeries	performent of the best of	n and the
+ etat	a tiger to taget to tage to taget taget to taget ta	ther the base there	·
etal_geon	1 12214031 10212142 5/195/30 1021244 5/195/30 10244 5/195/30 1044499 (uii) (uii ng) (beampest	pull HYDE,	88-K 2011-09-28
> Econter			
a the Taxana			
a the second			
+ mattat.goinn			
+ Charge			
+ interact			
* -@rtgar			
<ul> <li>-@-tger_data</li> </ul>			
<ul> <li>Is fopology</li> </ul>			
<ul> <li>Im postgrep</li> </ul>			
> Entertrate Troutin			
<ul> <li>An important intervention</li> </ul>			
Statistication			
h in the second second			

As we can see in Image 11, the minimum date is **1951-04-30** 

**Create geometry column in messstelle table:** In this section, we want to create a geometry column from **e32** and **n32** columns from the messstelle table. With the below code, we are able to create a new column and we set the name as **geom** 

```
ALTER TABLE hygrisc.messstelle ADD COLUMN geom geometry(Point, 25832);
UPDATE hygrisc.messstelle SET geom = ST_SetSRID(ST_MakePoint(e32, n32),
```

#### 25832);

5	a Dashbo	sard Properti	ies SQL Stat	istics Depende	ncies Dependents	<ul> <li>ebwas</li> </ul>	/elwasi@ielwas *								
> @ public	8 9		E Q - 1	1 · 6 B	B- T - 1	ferito	~	4 0		4	* # 18*				
✓ ③ sina	45 et	an/olwaniteh	was v	-	No. of Concession, Name		-		and the second second						
> []‡ Collations	Dury F	ditor Duiry	History									50	vatch Pad		
> No ETS Configurations				Cond 1											
<ul> <li>In FTS Determines</li> </ul>	1	elect + fro	C STOR. DESS.	tert ter											
As FTS Parsers															
> EFTS Templates															
> I Foreign Tables															
<ul> <li>R) Functions</li> </ul>															
Materialized Views															
> it periodelates															
w El Tables (6)	i.														
> If ow stations															
> mikatalog_ge															
> iii katalog_ge > iii katalog_stoff	Data D	enter et antitale		Marcal Contracts											
<ul> <li>Matalog_pe</li> <li>Matalog_stoff</li> <li>Matalog_stoff</li> </ul>	Data Or	utput Explain	Messages	Notifications										1	
katalog_pe     katalog_stoff     messurements     messurements	Data Da	Mortaenge.cm	Messages	Notifications	eusbeuðurchmesser, mm double precision		Natoriacher, ruhe, wap double precision		embeutaengecm double precision		obeikante, filter, om	•	unterkente, filter, om double precision	4	geom 🔺
Atalog_pe     Matalog_te     Modeling_stoff     Modeling_stoff     Modeling_stoff     Modeling_stoff     Modeling_stoff     Modeling_stoff     Modeling_Functions	Data Or ster	fiberlaenge.cm double precisio	Messages a sumptrot double pr	Notifications vterge_cm ecision	eusbeudurchmesser.,mm double precision	a bull	historischer,ruha,.eop double precision	a but	einbiaulaengecm double precision	<b>A</b> (5-0)	obeikante, filter, om double precision		unterkante, litter, cm double precision	A	geometry 🖨 🂽 01011000020E864
Atabig_pe     Atabig_tet     Adaig_tet     Mataig_tet     Mat	Data Or sbor	filterfaenge_cm double precisio	Messages a sumptrol (null 1000	Notifications viewge.cm ecision jout jout	euthautantmeaser,mm double precision	(null) 115	historischer, ruhe, wsp double precision	a put but	einbaulaenge, om double precision	1000 2000	obehavis, filter, on double precision	(nul) 10349	unterkante, filter, cm double precision	4 (huit) 9349	geometry a Contraction of the Co
> ⇒ katalog_pe     > ⇒ katalog_th     // measurements     > ⇒ measwert     > ₩ torsware     // Tragger Functions     > ₩ torsware     // Tragger Functions	Data Di ster	fiberta Explain fibertaenge_cm double precisio	Messages	Notifications visionge_ens inut just just just	eubeuturthmeser,mm doutse precision	(null) (null) (null)	Batoriacher, ruhe, wap double precision	A Jud Jud Jud	embaulaengecm double precision	2000 [mal]	oberkante, filter, om double precision	(null) 10349 (null)	unterkante, filter, om double precision	4 (hull) 9349 (hull)	geom geometry 🔒 💽 0101000020E864 0101000020E864 010100002028864
S Transformer	Data Or siter	dput Explain fiterisenge.cm double precisio	Messages Mes	Notifications visonge_ens ectrion jour jour jour jour jour jour jour jour	scabaudardhmesair, mm double precision	(nul) (nul) (nul) (nul)	historischer, nåre, veg double precision		einbaufaenge,.cm double precision	(pol) 2000 (pul) 635	oberkante, filter, om double precision	(null) 10349 (null) 7505	unterkante, Biter, om double precision	(hull) 9349 (hull) 7505	geometry 0101000020EB6- 0101000020EB6- 0101000020EB6- 0101000020EB6- 0101000020EB6-
<ul> <li>Transformer ()</li> </ul>	Data Or siber A RV media. 1 Pasetta. 1 Dataria.	Aput Explain Ateriaenge, om double precisio	Messages Mussinged Mussing	Notifications receiver init init init init init init	ausbeudurchmesserumm double precision	(rul) (rul) (rul) (rul) (rul)	historischer, ruhe, wep double precision	A Red Red Red Red Red	embaulaengecm double precision	(1000) (1000) (1000) (1000) (1000) (1000)	oberkante, filter, om double precision	(null) 10349 (null) 7505 (null)	unterkante, Biter, om dosble precision	(hul) 9349 (hul) 7505 (hul)	geom geometry & E 0101000020E864 0101000020E864 0101000020E864 0101000020E864
<ul> <li>Tratalog_pet</li> <li>Tratalog_pet</li> <li>Trassurements</li> <li>Trassurements</li> <li>Trassurements</li> <li>Trape Functions</li> <li>Trippet</li> <li>Trape functions</li> </ul>	Data Or siber A R4 media. 1 Padetta. 1 Debru. 3 Geteral. 1 Dasail.	ofput Explain Merlaenge.om double precisio	Messages bull full full full full full full full	Notifications viewge_en existen juit juit juit juit juit juit juit juit	eutbudachmesser, mm double precision	(rul) (rul) (rul) (rul) (rul) (rul)	hatorlacher, ruhe, wep double precision		erebudaenga.cm double precision	2000 (nul) 635 (nul) 900	oberkante, filter, om double precision	(null) 10349 (null) 7505 (null) 3452	unterkante, filter, om dosble precision	(hull) 9349 (hull) 7505 (hull) 3452	georety georety 0101000020E64 0101000020E64 0101000020E64 0101000020E64 0101000020E64 0101000020E64
S = Natalog_pe     S = Natalog_stoff     S = messatele     S = messatele     S = messatele     Wg Trigger Functions     Wg Trigger Functions     Wg Trigger Summary     Werwar (7)     S = moster     S = most     Columns     S = Columns     S = Columns	Data Or ster A R4 media. 3 Padetta. 1 Datau. 3 Gueteral. 1 Diazael.	ofput Explain Meriaesge.om double precisio	Messages Messages Messages Messages double pr built public publ	Notifications teergis.em put ind put put put put put put put put	eusbeudurchmesser.jom double perclann	(md) (md) (md) (md) (md) (md) (md)	historiacher, nufer, weip double precision		embudaenge.cm double precision	(mail) 2000 (mail) 635 (mail) 900 815	oberkants, filter, om double precision	(null) 10349 (null) 7505 (null) 3452 3150	unterkante, Biter, om double precision	(hul) 9349 (hul) 7505 (hul) 3452 3100	geometry (1010000205564 01010000205564 010100002055864 010100002055864 010100002055864 010100002055864 010100002055864
S = Natalog, pe     S = Natalog, pe     S = Natalog, stoff     S = measurements     S = measurement     S = measurement	Duto Di iber A Nimedia. 1 Pasetti, 1 Datauti, 1 Datauti, 1 Datauti,	fiterlaenge, om double precisio	Messages and Messages (0.00)	Notifications recision put indi put put put put put put put put	eutoudachmesermm doutie predision	(mail) (mail) (mail) (mail) (mail) (mail) (mail) (mail) (mail)	hatolischer, nähr, wip double precision	A Bori Jori Dori Dori Dori Dori Dori Dori	erebudaengaem double precision	(mal) (mal)	oberkante, filter, cm double precision	(null) 10349 (null) 7505 (null) 3462 3190 3668	unterkante, filter, enn doeble precision	(hull) 5349 (hull) 7505 (hull) 3452 3100 3068	geometry 0 0101000020286- 0101000020286- 0101000020286- 0101000020286- 0101000020286- 0101000020286- 0101000020286- 0101000020286-

The "messstelle" table has been enhanced with an additional column called "geom," which contains the geometry information representing the location of each station.

#### Merge two tables:

In this section, we aim to merge the "messwert" and "messstelle" tables based on the common column, "messstelle\_id." To achieve this, we will select the desired columns from each table and then perform the merge based on the "messstelle\_id" column.

```
select t1."messstelle_id", t1."name", t1.geom, t2."stoff_nr",
t2."messergebnis_c", t2."masseinheit",
t2."datum_pn", t2."messergebnis_cm" from hygrisc.messstelle t1 ,
hygrisc.nitrat t2
where t1."messstelle_id" = t2."messstelle_id";
```

ore	5 III 1. C	Q, Destoord	Properties	501. 3	Ratefics Depend	senten Dep	mdents # almost al	era di kan			
	* @ puble		a - 10	Q, ~	8 . 5 8	B'~ 1	+ isoni -		* * = *	5.5 4-	A 18-
	· · · · · · · · · · · · · · · · · · ·	9 dentate	STATISTICS.	25							
	* El Collatione	Curry Editor	Query Hitsto	ev.							Scratch Pad
	<ul> <li>B F75 Conferences</li> </ul>		-			-					
	5 Br. F.73 Dictoriantes	1	the set of	THESSE 7	secondaria con i	from alas.	callenterine + ta	dealer and the second	bound's." er-	stander treats."	
	<ul> <li>du F75 Painera</li> </ul>	1 ators	Th. Trainer	atta to	f" = t2, "manant	wite dd":	session of the		1.14		
	+ CFTS Templates			10125							
	<ul> <li>Emreige Tables</li> </ul>	a. 1									
	<ul> <li>Int Paradition</li> </ul>										
	F RAsteriolisted Wewes .										
	<ul> <li>Procetarez</li> </ul>										
	<ul> <li>Consequences</li> </ul>										
	<ul> <li>CLINERLINE</li> <li>XIII de déforte</li> </ul>										
	* IT hat shok on										
	+ minatalog.coott										
	* Etrossavorsetta	Data Oldput	Explan V	lescages	Nethanione						
	· Ittracesson		a Malates		anon	107.00 0	PROMIPORT A	massaicheit	a (mm.m	· messengebria.cm	
	* El retenent			161 -	Sound -	mpre -	South process -	101		- 164	
	* WTropper Fuections		200002119	PORTAL.	OPer pour incos+i.	104	100.249	1404	1800.09.01		
	* Types	1.1	90802091	TRAFIN.	OTEL BOOK1983840.		42,0969	rept.	1892-09-38	-	
	A second of the station on all		80302051	estade.	CONTRACTOR NAME	- 124	10,000	rege	10040011		
	5 Strated	A	10000101	THUS.	OF OT BIOCODEREM.	- 138	88.17	THE .	1440-00-08	- 55	
	* Stedart geom	- 3	80302178	STROE_	OF BET BUCKESSED BAD.	128	44.37	orbi	1400-07-15	-	
	* B Colorres		10002178	STERE_	01010002285840	134	-40.0940	1901	189440-19	-	
	> INTRONES	· · ·	290008113	Hall Han	of of Brospagse40.	104	78.3014	1991	1808-09-15		
	> 2+ Trooms		#1100x0	WD05	010130082083640.	104	10.81234	1401	180413.38	of Successfully 1	sar. Total-query nantime: 1 seco 480 mare: 73899 issue affected
	the second se		approximation of the	407101	CONTRACTOR DESIGNATION	1744	in the second se	stated in	1.100 million of the state.		

In order to have the above SQL command available as a new view for the subsequent section in QGIS, we should store it accordingly.

```
create view hygrisc.nitrat_geom as (select t1."messstelle_id", t1."name",
t1.geom, t2."stoff_nr", t2."messergebnis_c", t2."masseinheit",
t2."datum_pn", t2."messergebnis_cm" from hygrisc.messstelle t1 ,
hygrisc.nitrat t2
where t1."messstelle_id" = t2."messstelle_id")
```

# 2.5 QGIS

QGIS is a free and open-source geographic information system (GIS) software. It provides a wide range of tools and functionalities for visualizing, analyzing, and managing geospatial data. QGIS supports various data formats and allows users to create, edit, and publish maps.

You can download QGIS for free from the below link.

#### https://qgis.org/en/site/

The below video shows how to download and install QGIS for Windows which is highly recommended to watch before installing it.



To gain a better understanding of QGIS, I recommend watching the following video, which provides valuable insights and guidance on using the software.



**Create a time series video:** In this section, our objective is to generate a time series video depicting the changes in nitrate concentration over time in North Rhine-Westphalia, the most populous state in Germany. To begin, we must download the shapefile for North Rhine-Westphalia and import it into QGIS for further analysis and visualization.

Three below shapefiles need to be downloaded

- entire state shapefile (dvg1bld\_nw.shp)
- kreis shapefile (dvg1krs\_nw.shp)
- Gemeinde shapefile (dvg1gem\_nw.shp)

All three shapefiles can be downloaded from here. After downloading the shapefiles, we can proceed to load them into QGIS for visualization and analysis. The below video shows how to load the shapefile in QGIS.



Now we can see the map of NRW, kreis and Gemeinde. There are two options to create a video for time series.

**Locally with shapefile:** In here, we need to have a shapefile that consists of the nitrate concentration over time. download the notebook from here and run the Python codes to create two shapefiles. then we should load these two shapefiles to the QGIS. The first shapefile consists of all stations in NRW and the second one consists of the nitrate concentration.

The below video shows how we can load shapefiles to QGIS.



**Connect to Database:** The below video shows how we can connect our QGIS to Database and load the file from Database.



# **3 Dashboard**

In this section, the process of creating an interactive dashboard for our data will be explored. An interactive dashboard is a versatile tool that allows data to be interacted with, analyzed, visualized, and key information to be monitored by users.

This section discusses two approaches to creating a dashboard. The objective of this dashboard is to provide a user-friendly and interactive interface for data exploration and visualization, accessible even to non-programmers. Such a dashboard plays a crucial role in enhancing data comprehension and is widely utilized by managers and decision-makers to facilitate informed decision-making processes.

One notable example of this type of dashboard is the Covid-19 dashboard, which has gained widespread usage worldwide, including in Germany. The Covid-19 dashboard provides users with valuable insights into the number of new cases and deaths reported over various time periods. It helps individuals track the progression of the pandemic and understand the impact it has had on different regions and countries.

In our specific case, the objective is to develop a straightforward dashboard that showcases the map of North Rhine-Westphalia (NRW) alongside the concentration rates of Nitrate and Sulfate at different time intervals. This dashboard will provide a visual representation of the spatial distribution of these pollutants and enable users to observe any temporal variations in their concentrations within NRW.

### **Plotly Dash:**

**Plotly:** Plotly is a computing company located in Montreal, Canada. They develop online data analytics and visualization tools. Plotly offers online graphing, analytics, and statistics tools for their users, as well as scientific graphing libraries for Python, R, MATLAB, Perl, Julia, Arduino, and REST. Plotly offers several open-source and enterprise products such as Dash which have been used for creating simple and interactive dashboards in this project.

**Dash:** Dash is a framework to build data apps rapidly not only in Python but also in R, Julia, and F#. According to Plotly's official website, Dash is downloaded 800,000 times per month which shows that nowadays Dash getting more popular. Dash is a great framework for anyone who uses data with a customised user interface. Through a couple of simple patterns, Dash eliminated all of the technologies as well as protocols that are needed to make a full-stack web app with interactive data considerations. Another good feature is that Dash is running on web browsers so it means that no other application needs to run it.

To learn more about creating a dashboard with Plotly Dash, you can follow the link provided below. This resource contains comprehensive tutorials that guide you through the process of building a simple dashboard using Plotly Dash. These tutorials will provide you with step-by-step instructions and examples to help you create interactive and visually appealing dashboards using Plotly Dash.

### https://www.youtube.com/c/CharmingData

Dash is also offering some dashboards examples which could be really nice and helpful to get ideas. | Click here for Dash gallery

All the source codes of the dash gallery are available in | here

**Dashboard Design:** I have developed a web application dashboard that effectively visualizes time series data for Nitrate and Sulfate. The image below shows the main page of our dashboard.



In order to run the above dashboard on your local system, you need to do the following:

#### Click here and pull the repository

direct to the "Dashboard" folder, then all the Python codes and SQL queries are available to run the application.

There is a yml file inside the folder which name is "dashboard\_environment". This yml file will create an environment with all the necessary packages which you will need to run the application. The first step needs to load this file to the Anaconda, if you are not sure how to do that please refer to the 2.3.3 Anaconda section.

The next step is to add the credentials (Username, DB name, password and ...) of the database to the "credential\_temp" file.

And the last step is to run the "app.py" inside the new environment that you have already created with the help of "dashboard\_environment.yml"

# 4. Result

### Nitrate concentration 2000-2010

The video below shows the concentration of nitrate in North Rhine-Westphalia (NRW) from 2000 to 2010. This visualization was created using QGIS 3.16. By watching the video, you can observe the temporal changes in nitrate levels across NRW during the specified time period.



### Nitrate concentration 2010-2020

The video below shows the concentration of nitrate in NRW from 2010 to 2020. The video was created with QGIS 3.16



### Sulfat concentration 2000-2010

The video below shows the concentration of sulfate in NRW from 2000 to 2010. The video was created with QGIS 3.16



### Sulfat concentration 2010-2020

The video below shows the concentration of sulfate in NRW from 2010 to 2020. The video is created with QGIS 3.16



### 5. Project codes

All the codes are available in the below link.

Click here for project codes

# **Weitere Infos**

• EOLab-Wiki-Seiten zum Thema Grundwasserdaten in NRW

From: https://wiki.eolab.de/ - HSRW EOLab Wiki

Permanent link: https://wiki.eolab.de/doku.php?id=eolab:openhype:start



Last update: 2024/03/15 17:21